J. Perinat. Med.
6 (1978) 89

Umbilical cord complications as a cause of perinatal morbidity and mortality

Samuel Bruce, L. Stanley James, Edward Bowe, Henry Rey, Haider Shamsi

Division of Perinatal Medicine, Departments of Obstetrics and Pediatrics, College of Physicians and Surgeons, Columbia University, New York

The fetal heart pattern of variable deceleration has been correlated with compression of the umbilical cord [5]. It is the most frequently occurring pattern, being seen in approximately 30% of labors [2, 9, 10, 11, 12, 13]. Changing the maternal position will sometimes modify or even abolish this pattern[5]. This has led to the general belief that variable deceleration is benign, particularly if the heart rate does not fall below 80/min. As a consequence, the potential for the development of serious problems later in labor may be overlooked. Because of this we have examined the various factors contributing to perinatal mortality and morbidity. This study has revealed that compression of the umbilical cord is the major single factor associated with depression at birth in our clinic.

1 Material

The patient population includes 8038 infants born at the Columbia-Presbyterian Medical Center, New York City, over a three-year period from January 1974 to December 1976. All the known antepartum complications and identifiable complications at delivery were noted and tabulated for each stillbirth, neonatal death and all surviving newborns with an APGAR score of 6 or less at 1 or 5 minutes over the three years under review. Those infants having an APGAR score of 6 or less at 1 and 5

Curriculum vitae

Dr. SAMUEL BRUCE was born in 1941 in Ghana. He graduated from the University of Bridgeport, Connecticut, in 1965 and the Mt. Sinai School of Medicine in 1971. He completed his training in obstetrics and gynecology at New York University in 1976 and in Perinatal Medicine at Columbia University, College of Physicians and Surgeons, New York in 1978.

visually identified either as a loop or loops around the neck or body or as a prolapsed cord.

The data for the total number of births, stillbirths, neonatal deaths and perinatal mortality are presented by year in Tab. I. The stillbirth rate has remained essentially unchanged for 1974, 1975 and 1976 respectively. In Tab. II are listed by year the percentages of newborn infants with an APGAR Score < 6 at 1 and 5 minutes. It can be seen that the number of infants considered morbid has remained the same for each of the three years under review.

Tab. I. General statistics

	1974	1975	1976
Total deliveries*	2,790	2,714	2,534
Still births*	7.9/M	7.4/M	8.3/M
Neonatal deaths*	8.7/M	8.2/M	6.0/M
Perinatal mortality*	16.6/M	15.5/M	14.3/M

* Over 1000 gms.

Tab. II. APGAR scores at 1 and 5 minutes

	1974	1975	1976
Number of deliveries*	2,790	2,714	2,534
1 Minute APGAR ≤ 6	11.1%	9.2%	11.6%
5 Minute APGAR ≤ 6	2.4%	2.1%	2.4%

* Over 1000 gms

2 Results

In Tab. III are listed the factors associated with stillbirth in the three years under review. It can be seen that Rh sensitization (14.5%), umbilical cord complications (14.5%), abruptio placenta (13.2%) and congenital malformation (6.8%) represent the leading factors associated with all stillbirths that occurred in the review period. Rh sensitization is a leading factor at our institution because it is a referral center for this disease. **Umbilical cord complications are as frequently associated with stillbirths during this three year review.** The category of "others" (15.8%) represents twelve other single factors added together. In twenty-four of the cases (31.6%), the factors associated with stillbirth were unknown.

Factors associated with neonatal deaths are presented in Tab. IV. It can be seen that prematurity is associated with over one third of the deaths, congenital anomalies with 14.6%, while other causes represent the sum of forty single factors. Umbilical cord complications represent 3.1% of the factors associated with neonatal deaths.

Tab. V lists the factors observed to be associated with depression at birth during this three year survey. Morbidity (depression at birth) is defined as an APGAR score of ≤ 6 at 1 or 5 minutes.

Tab. III. Factors associated with stillbirths*. 3 year survey of 8038 births.

Factors	No.	% of factors
Rh	11	14.5%
Cord	11	14.5%
Abruptio	10	13.2%
Cong. anomalies	5	6.6%
Toxemias	2	2.6%
Prematurity	1	1.3%
Unknown	24	31.6%
Other Factors**	12	15.8%
Total	76	100%

* Over 1000 gms

** Maternal diabetes, amnionitis, placenta previa, maternal trauma, anemia, unregistered mother, meconium stained fluid at delivery, advanced maternal age.

Tab. IV. Factors associated with neonatal deaths*. 3-year survey of 8038 births.**

Factors	No.	% of total factors in population
Prematurity ¹⁾	37	38.5%
Cong. anomalies	14	14.6%
Rh disease	11	11.5%
Abruptio placentae	3	3.1%
Meconium	3	3.1%
Group B-strep	3	3.1%
Umb. cord compl.	3	3.1%
Unregistered mother	2	2.1%
Diabetes mellitus	2	2.1%
Amnionitis	2	2.1%
Other causes ²⁾	16	16.7%
Total	96	100%

* Liveborn infants dying during first 28 days of life

** Over 1000 gms

¹⁾ Less than 2500 gms²⁾ Small for gestational age, fetal anemia, incompetent cervix, maternal hypertension, preeclampsia, premature rupture of membranes, maternal drug addiction, prolonged uptake of membranes.

Umbilical cord complications are seen to be most frequently associated with depression at birth, accounting for 16.4% of the factors while meconium stained amniotic fluid is the second most frequent factor. Together these two constitute almost a third of all the factors associated with depression at birth.

Tab. V. Factors associated with perinatal morbidity.*
A 3-year survey of 8038 births**.

<i>Factor</i>	<i>No.</i>	<i>% of total population of factors</i>
Umbilical cord	111	16.4%
Meconium	81	12.0%
Prematurity	44	6.5%
Breech or abn. presnt.	41	6.1%
CPD	30	4.4%
Precipitous labor	24	3.6%
Post maturity	23	3.4%
Unregistered preg.	17	2.5%
Multiple pregnancy (Twin B)	17	2.5%
Placental problems	14	2.0%
Rh disease	12	1.8%
Toxemia	12	1.8%
Amnionitis	11	1.6%
Difficult delivery (C/S)	6	0.9%
Other factors***	219	32.4%
Unknown causes	14	2%
Total population of factors	676	100%

* Apgar score ≤ 6 at 1 or 5 minutes

** > 1000 grams

*** S.G.A., maternal anemia, fetal macrosomia, maternal agitation, undiagnosed twins, maternal hypoxemia, maternal drug addiction, intubation of newborn, vacuum extraction, difficult c/section, maternal supine hypotension, prolonged second stage, mid forceps, maternal over-medication etc.

Discussion

Of all the factors recognized as being associated with depression at birth, this survey has identified that complications of the umbilical cord occur most frequently. Meconium stained amniotic fluid was the second most frequent factor. Together these two factors were associated with nearly one-third of the depressed newborns. Cord complication was also the second most common factor associated with stillbirth and the seventh with neonatal death. In all instances the umbilical cord was identified to be in an abnormal position at birth, either around the neck or body or prolapsed. Cord complications could also have been unrecognized in some of the cases of meconium, as well as in some of the "unknown" group and could therefore

have been an even more frequently associated factor with depression at birth than 16.4%.

Cord complications could lead to depression at birth in a number of ways. The most likely is from hypoxia through intermittent occlusion. In addition, experiments by MING-NENG YEH and associates [14] have demonstrated that cord occlusion can cause myocardial conduction defects in the fetal baboon; similar changes with cardiac arrest have also been observed by S. C. YEH and associates in the human fetus during labor [15]. This effect appears to be separate from that of hypoxic depression, since it can be prevented by the prior administration of atropine. A third possible effect could be metabolic. DANIEL and associates have shown that intermittent hypoxia from occlusion of the umbilical cord in the fetal lamb leads to high levels of vasopressin and loss of solute, particularly sodium, in the urine [3, 4]. Another possible cause is hypovolemia. During partial cord occlusion when the vein only is likely to be occluded, blood flowing to the placenta and the umbilical arteries would not return to the fetus and would remain sequestered in the placenta.

The pattern of variable deceleration of the fetal heart rate, observed during labor, has been found to correlate with occlusion of the umbilical cord; it is also the most common abnormal heart rate pattern occurring in approximately 30% of all labors [2, 9, 10, 11, 12, 13]. This should provide the clinician with a means of recognizing a potential problem prior to birth and acid base analysis of fetal blood should permit early detection of those cases which are developing hypoxia and acidosis [1, 6, 7, 8]. Unfortunately, this is not always helpful since the most severe effect of cord occlusion may not occur until the last 15–20 minutes of the second stage of labor as the fetus descends. At this time the mother is usually in the delivery room and monitoring has been stopped. Undoubtedly, a better means of determining which cases of variable deceleration of fetal heart rate are likely to be associated with a poor outcome is needed. Closer attention to the fetal heart rate and fetal acid base state in the latter stages of labor should be helpful, particularly in the final 15–20 minutes of labor which could be the most critical time for a fetus with a cord complication. It

would seem, however, that additional diagnostic criteria should also be sought. These could include identification of the exact position of the cord by ultrasound, continuous fetal PO_2 with the transcutaneous electrode, and examination of the actual

fetal electrocardiogram. At present we can offer no single solution to the problem which remains probably the most important cause of depression at birth.

Summary

Examination of various factors contributing to perinatal mortality and morbidity has revealed that compression of the umbilical cord is the major single factor associated with depression at birth.

The fetal heart pattern of variable deceleration which has been correlated with compression of the umbilical cord [5] is also the most frequently occurring pattern, being seen in approximately 30% of labors [2, 9, 10, 11, 12, 13]. Because of a general belief that variable deceleration is benign, particularly if the heart rate does not fall below 80/min, the potential for the development of serious problems later in labor may be overlooked.

From a patient population of 8038 infants born at the Columbia-Presbyterian Medical Center, New York City, 1974–1976, all the known antepartum complications and identifiable complications at delivery were noted and tabulated for each stillbirth, neonatal death and all surviving newborns with an Apgar of 6 or less at 1 or 5 minutes over the three years under review. Malposition of the umbilical cord was visually identified either as a loop or loops around the neck or body or as a prolapsed cord. Umbilical cord complications were a leading factor associated with stillbirths during this three year review, being present 14.5% of the time. The category of "others" (15.8%) represents twelve other single factors added together. In twentyfour of the cases (31.6%), the factors associated with stillbirth were unknown.

Prematurity represents over a third of all the factors

associated with neonatal deaths, congenital anomalies 14.6%, while other causes represent the sum of forty single factors. Umbilical cord complications represent 3.1% of the factors associated with neonatal deaths.

Umbilical cord complications were most frequently associated with depression at birth, accounting for 16.4% of the factors while meconium stained amniotic fluid is the second most frequent factor. Together these two constitute almost a third of all the factors associated with morbid newborns. Cord complications could also have been unrecognized in some of the cases of meconium, as well as in some of the "unknown" group and could therefore have been an even more frequently associated factor with depression at birth than 16.4%.

Cord complications could lead to depression at birth in a number of ways. The most likely is from hypoxia through intermittent occlusion. Other causes include myocardial conduction defects [14, 15], metabolic effects from high levels of vasopressin and loss of solute, particularly sodium, in the urine [3, 4], and blood loss into the placenta if the cord is only partially occluded.

Fetal monitoring may not always be helpful if the most severe effect of cord occlusion occurs during the final 15–20 minutes of the second stage of labor, as the fetus descends. This could be the most critical time for the fetus with a cord round the neck; at this time the mother is usually in the delivery room and monitoring has been stopped.

Keywords: Perinatal morbidity, perinatal mortality, stillbirth, umbilical cord complications.

Zusammenfassung

Perinatale Morbidität und Mortalität infolge Nabelschnurkomplikationen

Bei der Durchsicht verschiedener Faktoren, die die perinatale Mortalität und Morbidität beeinflussen, hat sich gezeigt, daß die Kompression der Nabelschnur der wichtigste einzelne Faktor für die Neugeborenenendepression ist.

Das Herzfrequenzmuster der variablen Dezeleration, welches auf Nabelschnurkompression zurückgeführt wird [5], ist zugleich das häufigste Dezelerationsmuster, das in ca. 30% sub partu beobachtet wird [2, 9, 10, 11, 12, 13]. Auf Grund der allgemeinen Annahme, daß variable Dezelerationen ungefährlich seien, besonders dann, wenn die Herzfrequenz nicht unter 80/min abfällt, kann die mögliche Gefährdung zu einem späteren Zeitpunkt der Geburt übersehen werden.

Bei einer Population von 8083 am Columbia-Presbyterian Medical Center in New York City im Zeitraum von 1974–1976 geborenen Kindern wurden alle bekannten, ante-

partualen Komplikationen und erkennbaren Geburtskomplikationen registriert; jeder neonatale Todesfall und alle überlebenden Neugeborenen mit einer APGAR-Zahl < 6 nach 1 oder 5 Minuten wurden für die drei Jahrgänge tabelliert. Die Lage der Nabelschnur wurde mit dem Auge festgestellt und eingestuft entweder als ein- oder mehrmalige Umschlingungen um den Hals, den Körper oder als Nabelschnurvorfal.

Nabelschnurkomplikationen waren während dieser Dreijahresperiode in 14,5% vorhanden und damit ein wesentlicher Faktor in Zusammenhang mit Totgeburten. Die Kategorie „andere Faktoren“ (15,8%) bestand aus zwölf weiteren Einzelfaktoren, die zusammengefaßt wurden. In 24 Fällen (31,6%) waren die mit der Totgeburt einhergehenden Ursachen nicht bekannt.

Prämaturnität machte ein Drittel aller anderer Faktoren aus, die mit der neonatalen Sterblichkeit verknüpft waren, kongenitale Mißbildungen in 14,6%, während andere

Ursachen insgesamt 40 einzelne Faktoren beinhalteten. 3,1% der mit neonatalen Todesfällen einhergehenden Ursachen waren Nabelschnurkomplikationen.

Nabelschnurkomplikationen waren auch am häufigsten (in 16,4%) mit neonataler Depression verbunden, während mekoniumhaltiges Fruchtwasser das zweithäufigste Symptom war. Zusammen machen diese beiden Faktoren nahezu ein Drittel all jener Ursachen aus, die mit Neugeborenenmorbidity einhergehen. Nabelschnurumschlingungen könnten auch in manchen Fällen mit mekoniumhaltigem Fruchtwasser sowie in einigen Fällen aus der „unbekannten“ Ursachengruppe unerkannt geblieben sein und könnten demzufolge häufiger als in 16,4% Ursache einer Neugeborenenendepression sein.

Nabelschnurkomplikationen können auf vielerlei Weise zur Neugeborenenendepression führen. Der wahrschein-

lichste Mechanismus ist eine Hypoxie, bedingt durch intermittierenden Verschluss. Andere Ursachen sind: gestörte kardiale Überleitungszeit [14, 15], metabolische Einflüsse, bedingt durch hohe Spiegel an Vasopressin und Verlust von Salzen, insbesondere Natrium in den Urin [3, 4] sowie Blutpoolverschiebungen in die Plazenta dann, wenn die Nabelschnur nur partiell verschlossen wird.

Die fetale Herzfrequenzregistrierung ist nicht immer von Nutzen, vor allen Dingen dann nicht, wenn die schweren Kompressionssymptome während der letzten 15–20 Minuten der Austreibungszeit, also dann, wenn der Kopf tiefer tritt, auftreten. Dies dürfte der kritischste Zeitabschnitt für den Feten mit einer Nabelschnur um den Hals sein. Zu diesem Zeitpunkt liegt die Mutter meistens im Gebärsaal und die fetale Herzfrequenzüberwachung wurde beendet.

Schlüsselwörter: perinatale Morbidity, perinatale Mortality, Totgeburten, Nabelschnurkomplikationen

Résumé

Les complications du cordon ombilical: L'une des causes de la morbidité et de la mortalité périnatales

L'examen des diverses causes de la morbidité et de la mortalité périnatales a montré que la compression du cordon ombilical constitue le facteur isolé le plus important qui apparaisse dans l'état dépressif des nouveaux-nés.

Le modèle cardiaque foetal de décélération variable associée avec la compression du cordon ombilical [5] est le modèle le plus fréquent qu'on a pu relever, dans 30% environ des travaux d'accouchement [2, 9, 10, 11, 12, 13]. La décélération variable étant généralement considérée comme bénigne, surtout si la fréquence cardiaque ne baisse pas au-dessous de 80/min., on risque de négliger l'apparition éventuelle de problèmes graves en cours du travail.

Sur 8038 nouveaux-nés du Columbia-Presbyterian Medical Center, New York City, 1974–1976, nous avons enregistré toutes les complications intra-utérines connues et les complications identifiables à l'accouchement pour chaque cas de mort à la naissance ou néonatale et des nouveaux-nés ayant survécu avec un Apgar de 6 ou moins à 1 ou 5 minutes pendant les trois années de nos études. La mauvaise position du cordon ombilical est apparue sous forme d'une ou de plusieurs anses autour du cou ou du corps ou encore sous forme d'un cordon procident.

Les complications du cordon ombilical ont constitué l'un des facteurs principaux des morts à la naissance enregistrées pendant ces trois années d'observation, et cela dans un pourcentage de 14,5%. La catégorie des «autres» (15,8%) totalise 12 facteurs différents. Dans vingt-quatre des cas (31,6%), les facteurs associés avec les décès à la naissance sont restés inconnus.

La prématurité représente plus d'un tiers de tous les facteurs associés avec les morts néonatales, les anomalies congénitales 14,6% tandis que les autres causes totalisent quarante facteurs différents. Les complications du cordon ombilical représentent 3,1% des facteurs associés avec les morts néonatales.

Les complications du cordon ombilical constituent le plus un facteur de l'état dépressif à la naissance (16,4%), suivies par la coloration du liquide amniotique par le méconium. Ces deux facteurs représentent près d'un tiers de tous les facteurs observés chez les nouveaux-nés morbides. Il est possible qu'on n'ait pas toujours réussi à détecter des complications ombilicales dans certains cas de méconium ou de causes «inconnues» et que le pourcentage de ce facteur dans les états dépressifs des nouveaux-nés soit, en conséquence, supérieur encore à 16,4%.

Les complications ombilicales provoquent les états dépressifs à la naissance de diverses façons dont la plus probable est l'hypoxie par occlusion intermittente. Les autres causes comprennent les troubles de la conduction myocardiale [14, 15], les effets métaboliques de degrés élevés de vasopressine et la perte de solute, du sodium notamment, dans l'urine [3, 4], et la perte de sang dans le placenta s'il n'y a qu'occlusion partielle du cordon.

L'enregistrement du fœtus par monitor n'est pas toujours de grande utilité lorsque l'effet le plus grave de l'occlusion ombilicale survient durant les dernières 15–20 minutes de la seconde phase du travail, c.à.d. pendant la descente du fœtus. Ce pourrait être la période la plus critique pour le fœtus dont le cordon entoure le cou, car la mère est amenée en général à ce moment-là en salle d'accouchement après que l'enregistrement ait été stoppé.

Mots-clés: Complications du cordon ombilical, morbidité périnatale, mortalité périnatale, mort à la naissance.

Acknowledgement

This research has been supported by NIH Grants HD 07017, HL 14218 and GM 09069.

Bibliography

- [1] BOWE, E. T., R. W. BEARD, M. FINSTER, P. J. POPPERS, K. ADAMSONS, L. S. JAMES: Reliability of fetal blood sampling. *Amer. J. Obstet. Gynec.* 107 (1970) 279
- [2] CHAN, W. H.: Intrapartum fetal monitoring: *Obstet. and Gynec.* 41 (1973) 7
- [3] DANIEL, S. S., M.-N. YEH, E. T. BOWE, A. FUKUNAGA, L. S. JAMES: Renal response of the lamb fetus to partial occlusion of the umbilical cord. *J. Pediat.* 87 (1975) 788
- [4] DANIEL, S. S., M. K. HUSAIN, J. MILLIEZ, R. I. STARK, M.-N. YEH, L. S. JAMES: Renal response of the fetal lamb to complete occlusion of the umbilical cord. *Amer. J. Obstet. Gynec.* (1978) (In press)
- [5] HON, E. H.: An atlas of fetal heart rate patterns. New Haven, Hart Press 1968
- [6] KUBLI, F. W., E. H. HON, H. TAKEMURA: Observations on heart rate and pH in the human fetus during labor. *Amer. J. Obstet. Gynec.* 104 (1969) 1190
- [7] NERGES, A. T., L. MANN, A. BHAKTHAVATH, R. WEISS: Correlation of fetal heart rate uterine contraction patterns with fetal scalp blood pH. *Obstet. and Gynec.* 46 (1975) 392
- [8] NERGES, A. T., L. MANN, M. SANGHAVI, A. BHAKTHAVATHSALAN, R. WEISS: The association of umbilical cord complication and variable deceleration with acid base findings. *Obstet. and Gynec.* 49 (1977) 159
- [9] OTT, W. J.: Current status of intrapartum fetal monitoring. *Obstet. and Gynec.* 31 (1976) 339
- [10] PAUL, R. H., E. H. HON: Clinical fetal monitoring: V. Effect on perinatal outcome. *Amer. J. Obstet. Gynec.* 118 (1974) 529
- [11] REY, H. R., E. T. BOWE, L. S. JAMES: Impact of fetal heart rate monitoring and fetal blood sampling on infant mortality and morbidity. *Pediat. Res.* 8 (1974) 176
- [12] SHENKER, L.: Clinical experience with fetal heart rate monitoring of one thousand patients in labor. *Amer. J. Obstet. Gynec.* 115 (1973) 111
- [13] SHENKER, L., R. C. POST, J. S. SEILER: Routine electronic monitoring of fetal heart rate and uterine activity during labor. *Obstet. Gynec.* 46 (1975) 185
- [14] YEH, M.-N., H. O. MORISHIMA, W. H. NIEMANN, L. S. JAMES: Myocardial conduction defects in association with compression of the umbilical cord. *Amer. J. Obstet. Gynec.* 121 (1975) 951
- [15] YEH, S.-Y., B. ZANINI, R. H. PETRIE, E. H. HON: Intrapartum fetal cardiac arrest. *Obstet. and Gynec.* 50 (1977) 571

Received and accepted February 1, 1978

S. Bruce, M.D.
Dept. of Obstetrics and Gynecology
Columbia-Presbyterian Medical Center
New York City, N.Y. 10032 USA.