

Synthesis and Reactivity of Iridium(I) Fluorido Complexes: Oxidative Addition of SF₄ at *trans*-[Ir(F)(CO)(PEt₃)₂]

Nils Pfister,^[a] Minh Bui,^[a] Thomas Braun,^{*[a]} Philipp Wittwer,^[a] and Mike Ahrens^[a]

Dedicated to Prof. Manfred Scheer on the Occasion of his 65th Birthday

Abstract. The facile access to the Vaska type fluorido complexes *trans*-[Ir(F)(CO)(PR₃)₂] [**6**: R = Et, **7**: R = Ph, **8**: R = *i*Pr, **9**: R = Cy, **10**: R = *t*Bu] was achieved by halide exchange at *trans*-[Ir(Cl)(CO)(PR₃)₂] (**1–5**) with Me₄NF. Furthermore, the reaction of complex **6** with SF₄ gave *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt₃)₂] (**11**), whereas **8–10**

did not react. Reactivity studies revealed that **11** can selectively be manipulated at the sulfur atom by hydrolysis or fluoride abstraction to give *cis,trans*-[Ir(F)₂(SOF)(CO)(PEt₃)₂] (**12**) and *cis,trans*-[Ir(F)₂(SF₂)(CO)(PEt₃)₂][AsF₆] (**13**), respectively.

Introduction

Fluorinated transition metal complexes are highly interesting substances.^[1] Fluorido complexes of late transition metals can enable unique reaction pathways when compared to the reactivity of the corresponding metal complexes of the heavier halides.^[1a,1b,2] The preparation of transition metal fluorido complexes can be achieved either by treatment of metal fluorides with ligand precursors, by fluorination of suitable complexes or by the activation of element–fluorine bonds.^[1a,1b,21,3] Apart from C–F bond activation reactions,^[2e,2f,2h,21,4] this also includes reactions with SF₄ or SF₆.^[5] SF₄ is typically used as a tool for the fluorination of oxygen containing compounds, but has so far received only little attention in transition metal chemistry.^[5b,6] Replacement of a fluoride in SF₄ by nitrogen atoms or a bulky aryl group led to the discovery and commercialization of several deoxyfluorinating agents like FluoleadTM, DAST, MOST or Deoxo-Fluor[®].^[7] The derivatives often show a higher selectivity in fluorination, but they are not as versatile when compared with SF₄.^[6b,8]


Examples for transition metal derivatives of sulfur fluorides are scarce, but DFT calculations suggest the general feasibility of metal bound SF units.^[5e,5f,9] All experimentally available SF₃ complexes include only platinum group metal compounds.^[5b–5h] The most stable and best characterized derivatives are *trans*-[Ir(Cl)(F)(SF₃)(CO)(PEt₃)₂] and *trans*-[Pt(F)(SF₃)(PR₃)₂] (R = *i*Pr, Cy).^[5c–5h] The complexes

trans-[Pt(F)(SF₃)(PR₃)₂] (R = *i*Pr, Cy) were applied in deoxy-fluorination reactions of ethanol or benzophenone.^[5e,5h,10] Note that the reaction of SF₄ with **7** at 178 K to form a mixture of compounds with *cis,trans*-[Ir(F)₂(SF₃)(CO)(PPh₃)₂] as main product was also reported.^[5h] This demonstrates the influence of the fluorido ligand on the S–F bond cleavage in SF₄, because the chlorido complex **2** did give the respective SF₃ complex only at 323 K.^[5h] Other conversions at Vaska-type rhodium complexes were described as well, but the procedures always resulted in a mixture of products.^[5e] However, we recently reported on the selective formation of *cis,trans*-[Rh(F)₂(SF₃)(CO)(PEt₃)₂] by oxidative addition of SF₄ at *trans*-[Rh(F)(CO)(PEt₃)₂]. The complex was characterized by NMR spectroscopy and is only stable at low temperature in solution.^[5f]

In this contribution we report on the development of a reaction route to access iridium(I) fluorido carbonyl complexes. *trans*-[Ir(F)(CO)(PEt₃)₂] (**6**) is a useful starting compound for the synthesis of an iridium SF₃ complex, which can be handled in solution at room temperature. Reactivity studies towards hydrolysis and the Lewis acid AsF₅ are also presented.

Results and Discussion

Treatment of *trans*-[Ir(Cl)(CO)(PR₃)₂] with Me₄NF in dichloromethane gave *trans*-[Ir(F)(CO)(PR₃)₂] [**6**: R = Et, **7**: R = Ph, **8**: R = *i*Pr, **9**: R = Cy, **10**: R = *t*Bu] as the product of chlorine-fluorine exchange reactions (Scheme 1). Furthermore,



Scheme 1. Preparation of *trans*-[Ir(F)(CO)(PR₃)₂] [**6**: R = Et, **7**: R = Ph, **8**: R = *i*Pr, **9**: R = Cy, **10**: R = *t*Bu].

* Prof. Dr. T. Braun
E-Mail: thomas.braun@cms.hu-berlin.de

[a] Department of Chemistry
Humboldt Universität zu Berlin
Brook-Taylor-Straße 2
12489 Berlin, Germany

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/zaac.202000026> or from the author.

© 2020 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

the isotopologue of **6**, *trans*-[Ir(F)(¹³CO)(PEt₃)₂] (**6'**) was obtained when *trans*-[Ir(Cl)(¹³CO)(PR₃)₂] (**6'**) was treated with Me₄NF. The complex *trans*-[Ir(F)(CO)(PPh₃)₂] (**7**) has been described in the literature and was identified by its NMR and IR spectroscopic data (Table 1).^[11] Comparable fluorination reactions with Me₄NF in dichloromethane were reported before, for instance at rhodium and palladium complexes.^[2c,5f]

Table 1. Selected IR and NMR spectroscopic data for complexes *trans*-[Ir(F)(CO)(PR₃)₂] (**6–10**) (ν in cm⁻¹, δ in ppm, *J* in Hz).

Complex	ν(CO)	δ(¹⁹ F)	² J _{F,P}	δ(³¹ P)
6	1925	-265	33	25.6
7 ^[11]	1943	-254	30	24.4
8	1916	-256	26	49.5
9	1924	-255	26	39.3
10	1910	-243	24	71.5

Slow evaporation of a saturated *n*-hexane solution at room temperature produced crystals suitable for X-ray diffraction of **9** (Figure 1, left) and **10** (Figure 1, right). The CO and fluoro ligand in **9** are disordered. The asymmetric unit for complex **10** contains three molecules of which one reveals a disorder of the F, Ir and CO positions (Figure S31, Supporting Information). The bond lengths and angles found for all molecules of **10** are comparable, and hence only one is discussed in the following. The complexes *trans*-[Ir(F)(CO)(PCy₃)₂] (**9**) and *trans*-[Ir(F)(CO)(PtBu₃)₂] (**10**) exhibit an approximately square-planar coordination about the iridium center. Only a slight deviation from the ideal geometry is found for *trans*-[Ir(F)(CO)(PCy₃)₂] (**9**) with ligand–iridium–ligand angles of 88.37(2)° (F1–Ir1–P1) and 178.33(5)° (F1–Ir1–C1). A stronger distortion is observed for *trans*-[Ir(F)(CO)(PtBu₃)₂] (**10**) indicated by the F2–Ir2–C2 and P4–Ir2–P3 angles [169.88(13)° and 172.98(3)°], most probably due to the sterical demand of the *t*Bu groups at the phosphorus atoms. A nearly linear coordination of the carbonyl ligand to the metal center is revealed by the Ir–C–O angles of 176.3(12)° (**9**) and 176.4(3)° for (**10**) showing the expected bond lengths for the Ir–C [1.838(11) Å for **9**, 1.800(3) Å for **10**] as well as for the C–O [1.156(11) Å for **9**, 1.174(4) Å for **10**] distances.^[12] The metal carbon distance in **9** is longer than in *trans*-[Ir(Cl)(CO)(PCy₃)₂] (**4**)

[Ir–C: 1.788(10) Å],^[13] which is also observed when comparing the analogous rhodium complexes.^[5f,13–14] The Ir–F distances [1.965(5) Å for **9**, 1.9846(19) Å for **10**] are well comparable to these in other iridium(I) fluoro complexes as well as to the one observed in *trans*-[Ir(F)(CO)(P{C₆H₄-2-CH₃})₂].^[15]

The ¹⁹F NMR spectrum of the complex **6** reveals one signal at δ = -265 ppm which appears as a triplet (²J_{F,P} = 33 Hz) and can be assigned to an iridium(I) bound fluorine atom with two phosphorus nuclei in a *cis* position.^[11c,11d,12b,15a,15b,15e] The isotopologue *trans*-[Ir(F)(¹³CO)(PEt₃)₂] (**6'**) shows a doublet of triplets [²J_{C,F} = 84 Hz], caused by the coupling of the carbon atom of the carbonyl ligand to the fluorine atom in the *trans* position.^[5f,16] The ³¹P{¹H} NMR spectrum of compound **6** exhibits a single resonance at δ = 25.6 ppm showing the respective ²J_{F,P} doublet coupling. An additional doublet coupling can be detected for the ³¹P{¹H} NMR spectrum of the isotopologue **6'** with a ²J_{C,P} *cis*-coupling of 10 Hz.^[17] Compound **6'** reveals the expected doublet of triplet pattern in the ¹³C{¹H} NMR spectrum in the range of iridium(I) bound carbonyl ligands at δ = 174.1 ppm.^[17,18] A very intense absorption band at 1925 cm⁻¹ for the CO stretch mode can be detected in the IR spectrum of **6** (1878 cm⁻¹ for **6'**) which is well comparable to the known iridium based Vaska type complexes.^[12a,19] Compound **6** shows a significantly lower wavenumber than the compounds bearing heavier halogenido ligands.^[19] This phenomenon has already been discussed in the literature. It has been proposed that the fluoro ligand acts as a better π-donor than the other halogenido ligands.^[1a–1d] A push-pull interaction has also been reasoned for the stronger π backbonding from the metal to the ligand.^[2k,20] The NMR and IR spectroscopic data of the complexes **7–10** are very similar to those of **6** and are therefore not discussed (see Supporting Information and Table 1).

The fluoro complex **6** readily reacted with equimolar amounts of SF₄ in [D₈]THF, [D₈]toluene or CD₂Cl₂ below 263 K to give *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt₃)₂] (**11**) (Scheme 2). When introducing SF₄ to **6'** the formation of the isotopomer *cis,trans*-[Ir(F)₂(SF₃)(¹³CO)(PEt₃)₂] (**11'**) was observed. Attempts to isolate **11** resulted in the decomposition of





Figure 1. ORTEP diagram of *trans*-[Ir(F)(CO)(PCy₃)₂] (**9**) (left) and *trans*-[Ir(F)(CO)(PtBu₃)₂] (**10**) (right). Ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted for clarity. Selected bond lengths /Å and angles /° with estimated standard deviations in parentheses: **9**: Ir1–F1 1.965(5), Ir1–C1 1.838(11), Ir1–P1 2.3266(4), O1–C1 1.156(11); F1–Ir1–P1 88.37(2), F1–Ir1–C1 178.33(5), Ir1–C1–O1 178.99(11); **10**: Ir2–F2 1.9846(19), Ir2–C2 1.800(3), Ir2–P3 2.3810(9), Ir2–P4 2.3901(8), O2–C2 1.174(4), P3–Ir2–P4 172.98(3), P4–Ir2–F2 85.17(6), P4–Ir2–C2 93.96(10), F2–Ir2–C2 169.88(13), Ir2–C2–O2 176.4(3).

the complex giving SPEt_3 along with F_2PET_3 and unidentified iridium species.^[21] Based on ^{19}F NMR spectroscopy **11** was assumed to be one of four products in the reaction of *trans*- $[\text{Ir}(\text{NCS})(\text{CO})(\text{PET}_3)_2]$ with SF_4 , which was reported by Murdoch.^[5b] No oxidative addition of SF_4 was observed for **8–10** but the formation of traces of the respective difluorophosphoranes and phosphine sulfides were detected by ^{19}F and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy when the reaction mixtures were heated up to 323 K.^[21a,22] Comparable observations were reported for analogous rhodium compounds, and it was reasoned that the steric demand of the phosphine ligands plays a crucial role.^[5f]



Scheme 2. Formation of *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PET}_3)_2]$ (**11**).

DFT calculations were carried out on *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PR}_3)_2]$ (**11**; R = Et; R = *i*Pr) (see SI), and both complexes converged into stable structures. Furthermore, the calculated reaction enthalpies for the formation of *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PR}_3)_2]$ (**11**; R = Et; R = *i*Pr) from *trans*- $[\text{Ir}(\text{F})(\text{CO})(\text{PR}_3)_2]$ (**6**; R = Et, **8**; R = *i*Pr) and SF_4 both indicate the general feasibility of these reactions, whereas the formation of **11** was found to be more exergonic. However, as stated above, *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PiPr}_3)_2]$ was not observed experimentally. The analysis of the frontier orbitals of **11** reveals a HOMO, which is located at the sulfur atom and resembles an electron lone pair (Figure 2). The λ^4 -trifluorosulfanyl ligand should, therefore, be regarded as a one-electron donor.^[5e,5f,9g]




Figure 2. HOMO of the DFT-optimized structure of *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PET}_3)_2]$ (**11**). Hydrogen atoms at the phosphine ligands are omitted for clarity.

At 243 K the ^{19}F NMR spectrum of **11** shows four resonances at $\delta = 67.5$, -64.9 , -314.6 and -366.6 ppm which integrate in a ratio of 2:1:1:1 (Figure 3). The broad signal at $\delta = 67.5$ ppm can be assigned to the two fluorine atoms in the axial positions of the trigonal bipyramidal configuration about

the sulfur atom, and the resonance at -64.9 ppm to the one in the equatorial position.^[5d–5b] The signal for the sulfur bound fluorine atom at higher field shows a $^2J_{\text{F,F}}$ triplet splitting of 71 Hz which is in good accordance with fluorine-fluorine couplings in sulfur(IV) compounds.^[5c,5e,7a,7b,23] When cooling the sample to 203 K the resonance for the axial fluorine atoms at sulfur splits into two broad signals, indicating a dynamic behavior on the NMR timescale.




Figure 3. ^{19}F NMR spectrum (282.4 MHz) of *cis,trans*- $[\text{Ir}(\text{F})_2(\text{SF}_3)(\text{CO})(\text{PET}_3)_2]$ (**11**) at 243 K ($^\circ \text{SOF}_2$; * SF_6).

For the two metal fluoro ligands in **11** sharp doublets of triplets at $\delta = -314.6$ and -366.6 ppm appear in the ^{19}F NMR spectrum at a characteristic chemical shift.^[15a,24] Both show a $^2J_{\text{F,F}}$ coupling of 143 Hz indicating a mutual *cis* arrangement.^[15a,24b] Additionally the triplet couplings of 34 Hz and 29 Hz suggest that both metal fluorides have two phosphine ligands in a *cis* position.^[15a,24d] The ^{13}C labeled isotopologue **11'** doesn't show further splitting of the signals for the sulfur bound fluorine atoms, but an additional doublet splitting of 77 Hz and 5 Hz is observed for the resonances of the metal fluorides at $\delta = -314.6$ and -366.6 ppm, respectively.^[5f,16,24d–24f] A larger $^2J_{\text{F,C}}$ coupling of the carbonyl ligand to a fluoride in the *trans* position than to a fluoride in the *cis* position has been observed before, and this allows for an assignment of both resonances.^[5f,24d–24f] Thus, the signal at $\delta = -314.6$ ppm is attributed to the fluoro ligand *trans* to the carbonyl ligand and the resonance at $\delta = -366.6$ ppm to the fluoro ligand *cis* to the carbonyl ligand.^[5h,15a,24b,24e,24f]

At 203 K the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **11** shows one extremely broadened resonance at around $\delta \approx 13$ ppm (Figure 4). This broadening is presumably due to a hindered rotation of the λ^4 -trifluorosulfanyl group about the Ir–S bond. Similar fluxionalities have been observed for the above mentioned iridium, rhodium and platinum complexes.^[5c–5h] Increasing the temperature resulted at 298 K in a doublet of doublet pattern due to coupling to the two fluorine atoms in the *cis* position of the phosphines (Figure S15, Supporting Information). An additional doublet splitting of 7 Hz is detected in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **11'** at 298 K indicating a *cis* arrangement of the carbonyl ligand to the phosphine ligands. The above mentioned couplings are also observed in the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **11'** (Figure S19, Supporting Information), which shows a doublet of triplets of doublets in the expected range





Figure 4. Variable temperature $^{31}\text{P}\{^1\text{H}\}$ NMR spectra (121.5 MHz) of *cis,trans*-[Ir(F) $_2$ (SF $_3$)(CO)(PEt $_3$) $_2$] (**11**).

for CO ligands in iridium(III) carbonyl complexes at $\delta = 160.3$ ppm.^[24d,24f,25]

The trifluorosulfanyl complex **11** is very much prone to hydrolysis. Treatment of **11** with H $_2$ O in the presence of an excess of CsF resulted in the formation of *cis,trans*-[Ir(F) $_2$ (SOF)(CO)(PEt $_3$) $_2$] (**12**) (Scheme 3). The addition of CsF or Et $_3$ N is needed to remove the HF, which is formed by the hydrolysis. However, complex **12** converts further into a mixture of compounds within 2 days, of which **6**, SPEt $_3$ and F $_2$ PEt $_3$ were identified based on $^{31}\text{P}\{^1\text{H}\}$ and ^{19}F NMR spectroscopy (Figure S21, Supporting Information).^[21]



Scheme 3. Reactivity of *cis,trans*-[Ir(F) $_2$ (SF $_3$)(CO)(PEt $_3$) $_2$] (**11**).

The ^{19}F NMR spectrum of **12** shows three resonances at $\delta = 0.7$, -318.3 and -337.6 ppm of equal intensity. The broad multiplet at $\delta = 0.7$ ppm can be assigned to the fluorine atom of the SOF unit, which is in accordance with literature data.^[5e,5h] A doublet of triplets for one of the iridium bound fluorides is observed at $\delta = -318.3$ ppm exhibiting couplings to a fluoride as well as two phosphorus atoms in the *cis* position ($^2J_{\text{F,F}} = 140$, $^2J_{\text{F,P}} = 32$ Hz).^[5c,5f,15a,24b,24d–24f] For the other metal bound fluorine at $\delta = -337.6$ ppm a broad doublet with the $^2J_{\text{F,F}}$ of 140 Hz is detected.^[5f,15a,24b,24e]

In the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **12**, an AB pattern can be observed (Figure 5, bottom). The spectrum has been simulated (Figure 5, top) and the phosphine ligands couple to three fluorine atoms as well as the other phosphorus nucleus in the *trans*

position ($^2J_{\text{P,P}} = 342$ Hz).^[5c,5g,5h,26] The $^2J_{\text{P,F}}$ couplings appear in the typical range (A: 32, 28 Hz, B: 29, 33 Hz).^[5c,5f,15a,24d,24f] The $^3J_{\text{P,F}}$ couplings of the phosphine ligands to the sulfur bound fluorine atom of 7 Hz (A) and 10 Hz (B) are comparable to the ones found in *trans*-[Pt(F)(SOF)(PCy $_3$) $_2$] (**18**), *trans*-[Ir(Cl)(F)(SF $_3$)(CO)(PEt $_3$) $_2$] (**14**) and *trans*-[Pt(F)(SF $_3$)(PR $_3$) $_2$] (R = *i*Pr, Cy) (**15**).^[5c,5e]




Figure 5. Section of the simulated (top) and experimental (bottom) $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of *cis,trans*-[Ir(F) $_2$ (SOF)(CO)(PEt $_3$) $_2$] (**12**) at 298 K. The following shifts and coupling constants were obtained from the simulation: $\delta = 13.1$ (P $_a$, $^2J_{\text{P}_a,\text{P}_b} = 342.0$, $^2J_{\text{P}_a,\text{F}} = 27.6$, $^2J_{\text{P}_a,\text{F}} = 33.6$, $^3J_{\text{P}_a,\text{F}} = 5.5$ Hz) and 9.7 (P $_b$, $^2J_{\text{P}_a,\text{P}_b} = 342.0$, $^2J_{\text{P}_b,\text{F}} = 32.6$, $^2J_{\text{P}_b,\text{F}} = 28.6$, $^3J_{\text{P}_b,\text{F}} = 10.8$ Hz) ppm.

By addition of equimolar amounts of AsF $_5$ to **11** a sulfur bound fluoride of the SF $_3$ unit can be abstracted to give *cis,trans*-[Ir(F) $_2$ (SF $_2$)(CO)(PEt $_3$) $_2$][AsF $_6$] (**13**) along with unknown impurities (about 2% based on the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum; Figure S25, Supporting Information) (Scheme 3). Evaporation of the solvent resulted in a brownish oil as well as in an increasing amount of the side products. Similar reactions are reported for *trans*-[Ir(Cl)(F)(SF $_3$)(CO)(PEt $_3$) $_2$] and *trans*-[Pt(F)(SF $_3$)(PR $_3$) $_2$] (R = *i*Pr, Cy) yielding with BF $_3$ *trans*-[Ir(Cl)(F)(SF $_2$)(CO)(PEt $_3$) $_2$][BF $_4$] and *trans*-[Pt(F)(SF $_2$)(PR $_3$) $_2$][BF $_4$] (R = *i*Pr, Cy), respectively.^[5c,5e]

Four resonances were observed in the ^{19}F NMR spectrum of the SF $_2$ complex **13** at $\delta = -36.8$, -65.7 , -334.8 and -336.6 ppm in an integral ratio of 2:6:1:1 (Figure S26, Supporting Information). The most lowfield shifted resonance appears as a triplet of doublets ($^3J_{\text{F,P}} = 14$, $^3J_{\text{F,F}} = 11$, $^3J_{\text{F,F}} = 8$ Hz) and is in good accordance with a cationic SF $_2$ ligand bound to a transition metal center.^[5c,5e,5h] A broad signal at $\delta = -65.7$ ppm ($\Delta\nu_{1/2} = 1190$ Hz) appears for the anion AsF $_6^-$.^[27] The two resonances for the metal bound fluorides in **13** at $\delta = -334.8$ and -336.6 ppm show a typical $^2J_{\text{F,F}}$ coupling of 157 Hz accompanied by a strong roof effect for both doublet of triplet of triplets.^[5f,15a,24b,24e] The coupling to the equivalent phosphorus atoms of the phosphine ligands in the *cis* position results in a $^2J_{\text{P,P}}$ coupling constant of 32 Hz and 37 Hz, respectively.^[5c,5f,15a,24d,24f] A coupling to the two sulfur bound fluorine atoms is also observed in both resonances, exhibiting $^3J_{\text{F,F}}$ couplings of 11 Hz and 8 Hz. Comparable couplings have been

observed for other iridium(III) and platinum(II) complexes.^[5c,5e,24d,24f,28]

At 263 K, the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum shows a doublet of doublet of triplets at $\delta = 24.6$ ppm for **13** with corresponding coupling constants.^[5c,5e] The IR spectrum of the obtained oil shows prominent absorption bands at 692 cm^{-1} and 675 cm^{-1} , which can be attributed to modes of the hexafluoroarsenate anion (Figure S37, Supporting Information).^[27a,29] The CO stretching mode at 2064 cm^{-1} confirms the presence of an iridium compound in oxidation state +III.^[15a,24c,30]

Conclusions

In conclusion, we reported on the selective formation of *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (**11**) by reaction with SF₄. As starting compound *trans*-[Ir(F)(CO)(PR₃)₂] [**6**: R = Et, **7**: R = Ph, **8**: R = *i*Pr, **9**: R = Cy, **10**: R = *t*Bu] has to be prepared by exchange of the chlorido ligand. It is intriguing that only the latter complex **6** and *trans*-[Ir(F)(CO)(PPh₃)₂] react with SF₄ to yield the oxidative addition product, whereas *trans*-[Ir(F)(CO)(PR₃)₂] [**8**: R = *i*Pr, **9**: R = Cy, **10**: R = *t*Bu] do not react.

When compared to the analogous rhodium compound *cis,trans*-[Rh(F)₂(SF₃)(CO)(PEt)₂], complex **11** shows an enhanced stability allowing for manipulations at room temperature, although an isolation was not possible. SF₃ complexes are very rare, but the hydrolysis of **11** as well as the reaction with a Lewis acid is in accordance with the behavior of other transition metal or organo substituted λ^4 -trifluorosulfanyl units.^[5c,5e,7a,7c,31]

Experimental Section

General Methods and Instrumentation: The synthetic work was carried out with a Schlenk line or in a glove box in an atmosphere of argon. All reactions involving SF₄ and AsF₅ were performed on a stainless steel vacuum line with a defined volume. The reactions were performed in 15 mm o.d. and 3.8 mm o.d. PFA tubing.

Caution! Sulfur tetrafluoride and arsenic pentafluoride are highly toxic gases. Exposure to these chemicals can cause serious injuries.

All solvents were purified and dried by conventional methods and distilled under an atmosphere of argon before use or directly condensed into the reaction vessel. The complexes *trans*-[Ir(Cl)(CO)(PR₃)₂] [**1**: R = Et, **2**: R = Ph, **3**: R = *i*Pr, **4**: R = Cy] were prepared based on literature-known procedures. The preparation of *trans*-[Ir(Cl)(CO)(PrBu₃)₂] (**5**) was done analogous. SF₄ (99%), AsF₅ (99.9%) and Me₄NF were obtained from ABCR and used without further purification. Traces of thionyl fluoride and SF₆ were present in sulfur tetrafluoride but did not interfere with the chemistry. The NMR spectra were acquired with a Bruker DPX 300 or a Bruker Avance 300 spectrometer. The ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR chemical shifts were referenced to residual C₆D₅H at $\delta = 7.15$ ppm CDHCl₂ at $\delta = 5.32$ ppm or toluene-*d*₇ at $\delta = 2.09$ ppm. The $^{13}\text{C}\{^1\text{H}\}$ NMR chemical shifts were referenced to [D₈]toluene at $\delta = 137.84$ ppm. The ^{19}F NMR spectra were referenced to external CFCl₃ at $\delta = 0.0$ ppm and the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra to external 85% H₃PO₄ at $\delta = 0.0$ ppm. IR spectra were recorded with a Bruker Vertex 70 spectrometer that was equipped with

an ATR unit (diamond). Mass spectra were measured with a Micro-mass Q-TOF-2 mass spectrometer, which was equipped with a Linden LIFDI source (Linden CMS GmbH). Microanalyses were performed with a HEKAtech Euro EA Elemental Analyzer.

Structure Determination of Complexes 9 and 10: The data were collected at a BRUKER D8 VENTURE area detector, Mo- K_{α} radiation ($\lambda = 0.71073\text{ \AA}$). All measurements were performed at 100(2) K. Multi-scan absorption corrections implemented in SADABS were applied to the data.^[32] The structures were solved by intrinsic phasing method (SHELXT-2015) and refined by full-matrix least square procedures based on F^2 with all measured reflections (SHELXL-2015) with anisotropic temperature factors for all non-hydrogen atoms.^[33] All hydrogen atoms were added geometrically and refined on using a riding model.

Crystallographic data (excluding structure factors) for the structures in this paper have been deposited with the Cambridge Crystallographic Data Centre, CCDC, 12 Union Road, Cambridge CB21EZ, UK. Copies of the data can be obtained free of charge on quoting the depository numbers CCDC-1976836 (**9**) and CCDC-1976842 (**10**) (Fax: +44-1223-336-033; E-Mail: deposit@ccdc.cam.ac.uk, http://www.ccdc.cam.ac.uk).

General Procedure for the Synthesis of *trans*-[Ir(F)(CO)(PR₃)₂]: A PFA tube was loaded with Me₄NF (1.2 equiv.) and [Ir(Cl)(CO)(PR₃)₂] (**1–5**) (1 equiv.) dissolved in CH₂Cl₂ was added. After 15 min the solvent was removed in vacuo and the residue was extracted with *n*-hexane. Removing all volatiles from the extract gave **6** as a yellow oil and **7–10** as yellow powders in 98%, 80%, 97%, 82% and 93% yield.

Analytical Data for *trans*-[Ir(F)(CO)(PEt₃)₂] (6**):** ^1H NMR (300.1 MHz, C₆D₆, 298 K): $\delta = 1.81$ (mq, $^3J_{\text{H,H}} = 7.5$ Hz, 12 H, PCH₂CH₃), 1.06 ppm (mt, $^3J_{\text{H,H}} = 7.5$ Hz, 18 H, PCH₂CH₃); $^3J_{\text{H,H}}$ and $^3J_{\text{H,P}}$ -coupling constants were determined from a $^1\text{H}\{^{31}\text{P}\}$ NMR spectrum. ^{19}F NMR (282.4 MHz, C₆D₆, 298 K): $\delta = -264.8$ ppm (t, $^2J_{\text{F,P}} = 33$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, C₆D₆, 298 K): $\delta = 25.9$ ppm (d, $^2J_{\text{P,F}} = 33$ Hz) ppm. MS (LIFDI TOF, toluene): calcd. for C₁₃H₃₀FIrOP₂⁺ [M^+]: m/z 476.5, found: 476.2. IR (ATR, diamond): $\tilde{\nu} = 1925$ (vs, CO), 490 (m, Ir–F) cm^{-1} . Complex *trans*-[Ir(F)(¹³CO)(PEt₃)₂] (**6'**) was prepared in a similar manner by using *trans*-[Ir(Cl)(¹³CO)(PEt₃)₂] (**1'**). Selected spectroscopic data for **6'**: $^{13}\text{C}\{^1\text{H}\}$ NMR (75.5 MHz, [D₈]toluene, 298 K): $\delta = 174.1$ ppm (dt, $^2J_{\text{C,F}} = 84$, $^2J_{\text{C,P}} = 10$ Hz). ^{19}F NMR (282.4 MHz, C₆D₆, 298 K): $\delta = -264.8$ ppm (dt, $^2J_{\text{C,F}} = 84$, $^2J_{\text{F,P}} = 33$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, [D₈]toluene, 298 K): $\delta = 25.9$ ppm (dd, $^2J_{\text{P,F}} = 33$, $^2J_{\text{C,P}} = 10$ Hz). IR (ATR, diamond): $\tilde{\nu} = 1878$ (vs, ¹³CO) cm^{-1} .

Analytical Data for *trans*-[Ir(F)(CO)(PiPr₃)₂] (8**):** ^1H NMR (300.1 MHz, C₆D₆, 298 K): $\delta = 2.48$ [mhept, $^3J_{\text{H,H}} = 7.2$ Hz, 6 H, PCH(CH₃)₂], 1.31 ppm [md, $^3J_{\text{H,H}} = 7.2$ Hz, 36 H, PCH(CH₃)₂]; $^3J_{\text{H,H}}$ -coupling constants were determined from a $^1\text{H}\{^{31}\text{P}\}$ NMR spectrum. ^{19}F NMR (282.4 MHz, C₆D₆, 298 K): $\delta = -256.0$ ppm (t, $^2J_{\text{F,P}} = 26$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, C₆D₆, 298 K): $\delta = 49.45$ ppm (d, $^2J_{\text{P,F}} = 26$ Hz). MS (LIFDI TOF, toluene): calcd. for C₁₉H₄₂FIrOP₂⁺ [M^+]: m/z 560.2, found: 560.5. IR (ATR, diamond): $\tilde{\nu} = 1916$ (vs, CO), 487 (m, Ir–F) cm^{-1} . C₁₉H₄₂FIrOP₂ (560.23): calcd. C 40.77, H 7.56%; found C 41.15, H 7.64%.

Analytical Data for *trans*-[Ir(F)(CO)(PCy₃)₂] (9**):** ^1H NMR (300.1 MHz, C₆D₆, 298 K): $\delta = 2.43$ [m, 6 H, PCH(CH₂)₂(CH₂)₂CH₂], 2.19 [m, 12 H, PCH(CH₂)₂(CH₂)₂CH₂], 1.76 [m, 24 H, PCH(CH₂)₂(CH₂)₂CH₂], 1.26 ppm [m, 24 H, PCH(CH₂)₂(CH₂)₂CH₂]. ^{19}F NMR (282.4 MHz, [D₈]toluene, 298 K):

$\delta = -255.0$ ppm (t, $^2J_{\text{F,P}} = 26$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 39.3$ ppm (d, $^2J_{\text{P,F}} = 26$ Hz). **MS** (LIFDI TOF, toluene): calcd. for $\text{C}_{37}\text{H}_{66}\text{FIrOP}_2^+$ [M^+]: m/z 800.4, found: 800.7. **IR** (ATR, diamond): $\tilde{\nu} = 1924$ (vs, CO), 490 (m, Ir–F) cm^{-1} . $\text{C}_{37}\text{H}_{66}\text{FIrOP}_2$ (644.33): calcd. C 55.54, H 8.31; found C 55.41, H 8.27.

Analytical Data for *trans*-[Ir(F)(CO)(PrBu₃)₂] (10): ^1H NMR (300.1 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 1.60$ (pseudo triplet, $^3J_{\text{H,P}} - ^5J_{\text{H,P}} = 11.8$ Hz). ^{19}F NMR (282.4 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = -243.3$ ppm (t, $^2J_{\text{F,P}} = 24$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 71.5$ ppm (d, $^2J_{\text{P,F}} = 24$ Hz). **MS** (LIFDI TOF, toluene): calcd. for $\text{C}_{25}\text{H}_{54}\text{FIrOP}_2^+$ [M^+]: m/z 644.3, found: 644.6. **IR** (ATR, diamond): $\tilde{\nu} = 1910$ (vs, CO), 492 (m, Ir–F) cm^{-1} . $\text{C}_{25}\text{H}_{54}\text{FIrOP}_2$ (644.33): calcd. C 46.64, H 8.45; found C 46.78, H 8.54.

Formation of *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (11): In a PFA tube *trans*-[Ir(F)(CO)(PEt)₂] (**6**) (23 mg, 0.05 mmol) was dissolved in $[\text{D}_8]$ toluene. The solution was degassed in vacuo and 1 equiv. of SF₄ (0.05 mmol) was condensed into the PFA tube at 77 K. After sealing the PFA tube and storing the reaction mixture at 243 K for 16 h, *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (**11**) was identified by NMR spectroscopy. Selected spectroscopic data for **11**: ^1H NMR (300.1 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 1.94$ (mq, $^3J_{\text{H,H}} = 7.5$ Hz, 12 H, PCH₂CH₃), 0.96 ppm (mt, $^3J_{\text{H,H}} = 7.6$ Hz, 18 H, PCH₂CH₃); the $^3J_{\text{H,H}}$ -coupling constant was determined by a $^1\text{H}\{^{31}\text{P}\}$ NMR spectrum. ^{19}F NMR (282.4 MHz, $[\text{D}_8]$ toluene, 243 K): $\delta = 67.5$ (br., 2F, IrSF₂F), -64.9 (t, $^2J_{\text{F,F}} = 71$ Hz, 1F, IrSF₂F), -314.6 (dt, $^2J_{\text{F,F}} = 143$, $^2J_{\text{F,P}} = 34$ Hz, 1F, IrF), -366.6 ppm ($^2J_{\text{F,F}} = 143$, $^2J_{\text{F,P}} = 29$ Hz, 1F, IrF). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 11.7$ ppm (dd, $^2J_{\text{P,F}} = 34$, $^2J_{\text{P,P}} = 29$ Hz).

Complex *cis,trans*-[Ir(F)₂(SF₃)(¹³CO)(PEt)₂] (**11'**) was prepared in a similar manner by using *trans*-[Ir(F)(¹³CO)(PEt)₂] (**6'**). Selected spectroscopic data for **11'**: $^{13}\text{C}\{^1\text{H}\}$ NMR (75.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 160.33$ ppm (dtd, $^2J_{\text{C,F}} = 77$, $^2J_{\text{C,P}} = 7$, $^2J_{\text{C,F}} = 5$ Hz). ^{19}F NMR (282.4 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 67.5$ (br., 2F, IrSF₂F), -64.9 (br., 1F, IrSF₂F), -314.6 (dtd, $^2J_{\text{F,F}} = 143$, $^2J_{\text{F,P}} = 34$, $^2J_{\text{F,C}} = 77$ Hz, 1F, IrF), -366.6 ppm ($^2J_{\text{F,F}} = 143$, $^2J_{\text{F,P}} = 29$, $^2J_{\text{F,C}} = 5$ Hz, 1F, IrF). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 11.7$ ppm (ddd, $^2J_{\text{P,F}} = 34$, $^2J_{\text{P,P}} = 29$, $^2J_{\text{C,P}} = 7$ Hz). **MS** (LIFDI TOF, toluene): calcd. for $\text{C}_{12}^{13}\text{CH}_{30}\text{F}_4\text{IrOP}_2\text{S}^+$ [$\text{M}^+\text{-F}$]: m/z 566.11; found: m/z 566.13

Formation of *cis,trans*-[Ir(F)₂(SOF)(CO)(PEt)₂] (12): A sample of *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (**11**) (0.09 mmol) was prepared as described above, but either in the presence of 30 equivalents of CsF or 10 equivalents of Et₃N. The purity of the sample was verified by NMR spectroscopy at 243 K. The PFA tube was placed in a Young NMR tube, opened and 2 μL (1.2 Eq., 0.11 mmol) of H₂O was added with a microliter syringe at room temperature. The formation of **12** was monitored by NMR spectroscopy. Selected spectroscopic data for **12**: ^1H NMR (300.1 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 2.06$ –1.73 (m, 12 H, PCH₂CH₃), 0.96 ppm (m, 18 H, PCH₂CH₃). ^{19}F NMR (282.4 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 0.7$ (br. m, 1F, SOF), -318.3 (dt, $^2J_{\text{F,F}} = 140$, $^2J_{\text{F,P}} = 32.4$ Hz, 1F, IrF), -337.6 ppm (dm, $^2J_{\text{F,F}} = 140$ Hz, 1F, IrF). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, $[\text{D}_8]$ toluene, 298 K): $\delta = 11.3$ ppm (m, $^2J_{\text{P,P}} = 342$, $^2J_{\text{P,F}} = 33$, $^2J_{\text{P,F}} = 32$, $^2J_{\text{P,F}} = 29$, $^2J_{\text{P,F}} = 28$, $^3J_{\text{P,F}} = 10$, $^3J_{\text{P,F}} = 7$ Hz). The chemical shift and the coupling constants of the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum were obtained by simulation.

Treatment of *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (11) with AsF₅: A sample of *cis,trans*-[Ir(F)₂(SF₃)(CO)(PEt)₂] (**11**) (0.09 mmol) was prepared in CD₂Cl₂ leaving the reaction tube connected to the stainless

steel line. After a reaction time of 12 h at 263 K the mixture was degassed and equimolar amounts of AsF₅ were condensed onto the solution. The PFA tube was sealed and transferred to the NMR spectrometer. The formation of *cis,trans*-[Ir(F)₂(SF₂)(CO)(PEt)₂][AsF₆] (**13**) was observed by NMR spectroscopy upon thawing of the mixture. Spectroscopic data for **13**: ^1H NMR (300.1 MHz, CD₂Cl₂, 263 K): $\delta = 2.21$ (mq, $^3J_{\text{H,H}} = 7.7$ Hz, 12 H, PCH₂CH₃), 1.22 ppm (mt, $^3J_{\text{H,H}} = 7.7$ Hz, 18 H, PCH₂CH₃); the $^3J_{\text{H,H}}$ -coupling constant was determined by a $^1\text{H}\{^{31}\text{P}\}$ NMR spectrum. ^{19}F NMR (282.4 MHz, CD₂Cl₂, 263 K): $\delta = -36.8$ (tdd, 2F, SF₂), -65.7 (br., 6F, AsF₆), -334.8 (dtd, $^2J_{\text{F,F}} = 157$, $^2J_{\text{F,P}} = 32$, $^3J_{\text{F,P}} = 11$ Hz, 1F, IrF), -336.6 ppm (dtd, $^2J_{\text{F,F}} = 157$, $^2J_{\text{F,P}} = 37$, $^3J_{\text{F,F}} = 8$ Hz, 1F, IrF). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, CD₂Cl₂, 263 K): $\delta = 24.6$ ppm (dtd, $^2J_{\text{P,F}} = 37$, $^2J_{\text{P,P}} = 32$, $^3J_{\text{P,F}} = 14$ Hz). **IR** (ATR, diamond): $\tilde{\nu} = 2064$ (s, CO), 692 (vs, AsF₆), 675 (vs, AsF₆) cm^{-1} .

Supporting Information (see footnote on the first page of this article): Spectra, computational details, crystallographic data.

Acknowledgements

We acknowledge the DFG (Deutsche Forschungsgemeinschaft) for financial support. We also thank Stefan Sander for scientific discussions. Open access funding enabled and organized by Projekt DEAL.

Keywords: Sulfur fluorides; Iridium; S–F activation; Fluorido complexes; Sulfur tetrafluoride

References

- a) N. M. Doherty, N. W. Hoffmann, *Chem. Rev.* **1991**, *91*, 553–573; b) E. F. Murphy, R. Murugavel, H. W. Roesky, *Chem. Rev.* **1997**, *97*, 3425–3468; c) K. Fagnou, M. Lautens, *Angew. Chem.* **2002**, *114*, 26–49; *Angew. Chem. Int. Ed.* **2002**, *41*, 26–47; d) V. Grushin, *Chem. Eur. J.* **2002**, *8*, 1006–1014; e) R. Szpera, D. F. J. Moseley, L. B. Smith, A. J. Sterling, V. Gouverneur, *Angew. Chem.* **2019**, *131*, 14966–14991; *Angew. Chem. Int. Ed.* **2019**, *58*, 14824–14848; f) F. Nahra, M. Brill, A. Gómez-Herrera, C. S. J. Cazin, S. P. Nolan, *Coord. Chem. Rev.* **2016**, *307*, 65–80; g) M. G. Campbell, A. J. Hoover, T. Ritter, *Transition Metal-Mediated and Metal-Catalyzed Carbon–Fluorine Bond Formation*, in *Organometallic Fluorine Chemistry* (Eds.: T. Braun, R. P. Hughes), Springer International Publishing: Cham, **2015**, pp. 1–53; h) A. Vigalok, *Acc. Chem. Res.* **2015**, *48*, 238–247; i) A. Vigalok, *Organometallics* **2011**, *30*, 4802–4810.
- a) J. Miró, C. del Pozo, *Chem. Rev.* **2016**, *116*, 11924–11966; b) C. Berg, T. Braun, R. Laubenstein, B. Braun, *Chem. Commun.* **2016**, *52*, 3931–3934; c) J. Kohlmann, T. Braun, R. Laubenstein, R. Herrmann, *Chem. Eur. J.* **2017**, *23*, 12218–12232; d) J. Kohlmann, T. Braun, *J. Fluorine Chem.* **2017**, *203*, 140–145; e) M. F. Kuehnel, D. Lentz, T. Braun, *Angew. Chem.* **2013**, *125*, 3412–3433; *Angew. Chem. Int. Ed.* **2013**, *52*, 3328–3348; f) E. Clot, O. Eisenstein, N. Jasim, S. A. Macgregor, J. E. McGrady, R. N. Perutz, *Acc. Chem. Res.* **2011**, *44*, 333–348; g) V. V. Grushin, *Acc. Chem. Res.* **2010**, *43*, 160–171; h) H. Amii, K. Uneyama, *Chem. Rev.* **2009**, *109*, 2119–2183; i) V. V. Grushin, *Chem. Eur. J.* **2002**, *8*, 1006–1014; j) J. L. Kiplinger, T. G. Richmond, C. E. Osterberg, *Chem. Rev.* **1994**, *94*, 373–431; k) A. Mezzetti, C. Becker, *Helv. Chim. Acta* **2002**, *85*, 2686–2703; l) T. Braun, R. N. Perutz, *Chem. Commun.* **2002**, 2749–2757; m) J. M. Brown, V. Gouverneur, *Angew. Chem.* **2009**, *121*, 8762–8766; *Angew. Chem. Int. Ed.* **2009**, *46*, 8610–8614.
- a) N. Arnold, R. Bertermann, F. M. Bickelhaupt, H. Braunschweig, M. Drisch, M. Finze, F. Hupp, J. Poater, J. A. P.

- Sprenger, *Chem. Eur. J.* **2017**, *23*, 5948–5952; b) J. Bauer, H. Braunschweig, K. Kraft, K. Radacki, *Angew. Chem.* **2011**, *123*, 10641–10644; *Angew. Chem. Int. Ed.* **2011**, *50*, 10457–10460; c) M. A. Ellwanger, S. Steinhauer, P. Golz, H. Beckers, A. Wiesner, B. Braun-Cula, T. Braun, S. Riedel, *Chem. Eur. J.* **2017**, *23*, 13501–13509.
- [4] a) N. A. LaBerge, J. A. Love, *Activation and Formation of Aromatic C–F Bonds*, in *Organometallic Fluorine Chemistry* (Eds.: T. Braun, R. P. Hughes), Springer International Publishing: Cham, **2015**, pp. 55–111; b) O. Eisenstein, J. Milani, R. N. Perutz, *Chem. Rev.* **2017**, *117*, 8710–8753; c) T. Ahrens, J. Kohlmann, M. Ahrens, T. Braun, *Chem. Rev.* **2015**, *115*, 931–972; d) T. Fujita, K. Fuchibe, J. Ichikawa, *Angew. Chem.* **2019**, *131*, 369–408; *Angew. Chem. Int. Ed.* **2019**, *58*, 390–402; e) S. A. Macgregor, *Chem. Soc. Rev.* **2007**, *36*, 67–76; f) A. D. Sun, J. A. Love, *Dalton Trans.* **2010**, *39*, 10362–10374; g) W. Chen, C. Bakewell, M. R. Crimin, *Synthesis* **2017**, *49*, 810–821; h) S. A. Johnson, J. A. Hatnean, M. E. Doster, *Functionalization of Fluorinated Aromatics by Nickel-Mediated C–H and C–F Bond Oxidative Addition: Prospects for the Synthesis of Fluorine-Containing Pharmaceuticals*, in *Prog. Inorganic Chemistry* (Ed.: K. D. Karlin), Wiley: New Jersey, **2011**, Vol. 57, pp. 255–352; i) J. Weaver, S. Senaweera, *Tetrahedron* **2014**, *70*, 7413–7428; j) W. D. Jones, *Dalton Trans.* **2003**, 3991–3995; k) R. P. Hughes, *Eur. J. Inorg. Chem.* **2009**, 2009, 4591–4606; l) H. Torrens, *Coord. Chem. Rev.* **2005**, *249*, 1957–1985; m) Q. Shen, Y.-G. Huang, C. Liu, J.-C. Xiao, Q.-Y. Chen, Y. Guo, *J. Fluorine Chem.* **2015**, *179*, 14–22; n) P. Fischer, K. Götz, A. Eichhorn, U. Radius, *Organometallics* **2012**, *31*, 1374–1383; o) M. W. Kuntze-Fechner, C. Kerpen, D. Schmidt, M. Häring, U. Radius, *Eur. J. Inorg. Chem.* **2019**, 2019, 1767–1775; p) T. Schaub, P. Fischer, A. Steffen, T. Braun, U. Radius, A. Mix, *J. Am. Chem. Soc.* **2008**, *130*, 9304–9317.
- [5] a) M. Wozniak, T. Braun, M. Ahrens, B. Braun-Cula, P. Wittwer, R. Herrmann, R. Laubenstein, *Organometallics* **2018**, *37*, 821–828; b) T. Braun, D. Dirican, N. Pfister, M. Wozniak, *Chem. Eur. J.* **2019** DOI: 10.1002/chem.201904493; c) R. W. Cockman, E. A. V. Ebsworth, J. H. Holloway, *J. Am. Chem. Soc.* **1987**, *109*, 2194–2195; d) R. W. Cockman, E. A. V. Ebsworth, J. H. Holloway, H. M. Murdoch, N. Robertson, P. G. Watson, *Reaction of Nonmetal Fluorides with some Platinum Metal-Complexes*, in *Inorganic Fluorine Chemistry* (Ed.: J. S. Thrasher), ACS Symposium Series 555; American Chemical Soc: Washington DC, **1994**, pp. 326–337; e) C. Berg, T. Braun, M. Ahrens, P. Wittwer, R. Herrmann, *Angew. Chem.* **2017**, *129*, 4364–4368; *Angew. Chem. Int. Ed.* **2017**, *56*, 4300–4304; f) N. Pfister, T. Braun, P. Wittwer, M. Ahrens, *Z. Anorg. Allg. Chem.* **2018**, *644*, 1064–1070; g) P. Watson, *Reactions of the Sulphur Group Tetrafluorides with Some Rhodium Complexes*, Dissertation Thesis, University of Edinburgh, UK **1990**; h) H. M. Murdoch, NMR Studies of some novel Fluoro-Iridium-Complexes, Dissertation Thesis, University of Edinburgh, **1991**; i) L. Zámostná, T. Braun, B. Braun, *Angew. Chem.* **2014**, *126*, 2783–2787; *Angew. Chem. Int. Ed.* **2014**, *53*, 2745–2749; j) L. Zámostná, T. Braun, *Angew. Chem.* **2015**, *127*, 10798–10802; *Angew. Chem. Int. Ed.* **2015**, *54*, 10652–10656; k) T. A. McTeague, T. F. Jamison, *Angew. Chem.* **2016**, *128*, 15296–15299; *Angew. Chem. Int. Ed.* **2016**, *55*, 15072–15075.
- [6] a) M. R. C. Gerstenberger, A. Haas, *Angew. Chem.* **1981**, *93*, 659–680; *Angew. Chem. Int. Ed. Engl.* **1981**, *20*, 647–667; b) C. Ni, M. Hu, J. Hu, *Chem. Rev.* **2015**, *115*, 765–825; c) G. A. Boswell Jr., W. C. Ripka, R. M. Scribner, C. W. Tullock, *Fluorination by Sulfur Tetrafluoride*, in *Organic Reactions*; John Wiley & Sons, Inc., **1974**, Vol. 21, pp. 1–124; d) C.-L. J. Wang, *Fluorination by Sulfur Tetrafluoride*, in *Organic Reactions* (Ed.: A. S. Kende), John Wiley & Sons: Hoboken, NJ, **1985**, vol. 34, pp. 319–400; e) J. T. Goettel, D. Turnbull, M. Gerken, *J. Fluorine Chem.* **2015**, *174*, 8–13; f) A. L. Oppegard, W. C. Smith, E. L. Muetterties, V. A. Engelhardt, *J. Am. Chem. Soc.* **1960**, *82*, 3835–3838.
- [7] a) T. Umemoto, R. P. Singh, Y. Xu, N. Saito, *J. Am. Chem. Soc.* **2010**, *132*, 18199–18205; b) V. E. Pashinnik, E. G. Martyniuk, M. R. Tabachuk, Y. G. Shermolovich, L. M. Yagupolskii, *Synth. Commun.* **2003**, *33*, 2505–2509; c) V. E. Pashinnik, *J. Fluorine Chem.* **2002**, *117*, 85–98.
- [8] a) R. P. Singh, J. n. M. Shreeve, *Synthesis* **2002**, 2561–2578; b) L. N. Markovski, V. E. Pashinnik, *Synthesis* **1975**, 1975, 801–802; c) M. Hudlicky, *Fluorination with Diethylaminosulfur Trifluoride and Related Aminofluorosulfuranes*, in *Organic Reactions* (Ed.: A. S. Kende), John Wiley & Sons, **1988**, Vol. 35, pp. 513–637.
- [9] a) Y. Zhu, N. Li, R. Bruce King, *Inorg. Chim. Acta* **2019**, *486*, 332–339; b) X. Zeng, N. Li, R. B. King, *Polyhedron* **2019**, *163*, 33–41; c) H. Jiang, N. Li, R. Bruce King, *RSC Adv.* **2016**, *6*, 18874–18880; d) X. Gao, S. Gong, N. Li, R. B. King, *New J. Chem.* **2015**, *39*, 4939–4947; e) X. Gao, N. Li, R. B. King, *Inorg. Chem.* **2014**, *53*, 12635–12642; f) J. Deng, Q.-s. Li, Y. Xie, R. B. King, *J. Fluorine Chem.* **2013**, *146*, 37–45; g) J. Deng, C. Wang, Q.-s. Li, Y. Xie, R. B. King, H. F. Schaefer, *Inorg. Chem.* **2011**, *50*, 2824–2835.
- [10] C. Berg, N. Pfister, T. Braun, B. Braun-Cula, *Chem. Eur. J.* **2018**, *24*, 7985–7990.
- [11] a) C. A. McAuliffe, R. Pollock, *J. Organomet. Chem.* **1974**, *77*, 265–268; b) L. Vaska, J. Peone, E. R. Wonchoba, G. W. Parshall, *Fluoro Complexes of Rhodium(I) and Iridium(I)*, in *Inorganic Syntheses* (Ed.: G. W. Parshall), McGraw Hill: New York, **1974**, pp. 64–68; c) A. F. Williams, S. Bhaduri, A. G. Maddock, *J. Chem. Soc., Dalton Trans.* **1975**, 1958–1962; d) M. A. Cairns, K. R. Dixon, J. J. McFarland, *J. Chem. Soc., Dalton Trans.* **1975**, 1159–1164.
- [12] a) S. A. Cotton, *Rhodium and Iridium*, in *Chemistry of Precious Metals*; (Ed.: S. A. Cotton), Springer Netherlands: Dordrecht, **1997**, pp. 78–172; b) H. Werner, K. Ilg, B. Weberndörfer, *Organometallics* **2000**, *19*, 3145–3153.
- [13] E. Grobbelaar, S. Lötter, H. G. Visser, J. Conradie, W. Purcell, *Inorg. Chim. Acta* **2009**, *362*, 3949–3954.
- [14] a) E. Kuwabara, R. Bau, *Acta Crystallogr., Sect. C* **1994**, *50*, 1409–1411; b) M. R. Churchill, C. H. Lake, C. A. Miller, J. D. Atwood, *J. Chem. Crystallogr.* **1994**, *24*, 557–560.
- [15] a) J. Fawcett, D. A. J. Harding, E. G. Hope, *Dalton Trans.* **2010**, 39, 5827–5832; b) K. Ilg, H. Werner, *Organometallics* **1999**, *18*, 5426–5428; c) B. J. Truscott, F. Nahra, A. M. Z. Slawin, D. B. Cordes, S. P. Nolan, *Chem. Commun.* **2015**, *51*, 62–65; d) M. Gorol, N. C. Mösch-Zanetti, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, *Eur. J. Inorg. Chem.* **2004**, *2004*, 2678–2682; e) K. Ilg, H. Werner, *Organometallics* **2001**, *20*, 3782–3794.
- [16] a) B. T. Heaton, J. A. Iggo, C. Jacob, H. Blanchard, M. B. Hursthouse, I. Ghatak, M. E. Harman, R. G. Somerville, W. Heggie, P. R. Page, I. Villax, *J. Chem. Soc., Dalton Trans.* **1992**, 2533; b) J. Gil-Rubio, B. Weberndörfer, H. Werner, *J. Chem. Soc., Dalton Trans.* **1999**, 1437–1444.
- [17] a) H.-C. Böttcher, M. Junk, P. Mayer, W. Beck, *Eur. J. Inorg. Chem.* **2015**, *2015*, 3323–3327; b) D. Capitani, P. Mura, *Inorg. Chim. Acta* **1997**, *258*, 169–181.
- [18] H. A. Mayer, W. C. Kaska, *Chem. Ber.* **1995**, *128*, 95–98.
- [19] a) L. D. Field, E. T. Lawrenz, A. J. Ward, *Polyhedron* **1999**, *18*, 3031–3034; b) A. Tahara, Y. Miyamoto, R. Aoto, K. Shigeta, Y. Une, Y. Sunada, Y. Motoyama, H. Nagashima, *Organometallics* **2015**, *34*, 4895–4907.
- [20] J. P. Flemming, M. C. Pilon, O. Y. Borbulevitch, M. Y. Antipin, V. V. Grushin, *Inorg. Chim. Acta* **1998**, *280*, 87–98.
- [21] a) R. Bartsch, O. Stelzer, R. Schmutzler, *J. Fluorine Chem.* **1982**, *20*, 85–88; b) G. Baccolini, C. Boga, M. Mazzacurati, *J. Org. Chem.* **2005**, *70*, 4774–4777.
- [22] a) C. B. Caputo, D. Winkelhaus, R. Dobrovetsky, L. J. Hounjet, D. W. Stephan, *Dalton Trans.* **2015**, *44*, 12256–12264; b) M. Fild, R. Schmutzler, *J. Chem. Soc. A* **1970**, 2359–2364; c) A. Cleaves Peter, E. Kefalidis Christos, M. Gardner Benedict, F. Tuna, J. L. McInnes Eric, W. Lewis, L. Maron, T. Liddle Stephen, *Chem. Eur. J.* **2017**, *23*, 2950–2959; d) H.-U. Steinberger, B. Ziemer, M. Meisel, *Acta Crystallogr., Sect. C* **2001**, *57*, 835–837; e) W. I. Cross, S. M. Godfrey, S. L. Jackson, C. A. McAuliffe, R. G. Pritchard, *J. Chem. Soc., Dalton Trans.* **1999**,

- 2225–2230; f) C. F. Shaw, A. A. Isab, J. D. Hoeschele, M. Starich, J. Locke, P. Schulteis, J. Xiao, *J. Am. Chem. Soc.* **1994**, *116*, 2254–2260.
- [23] W. Gombler, *J. Fluorine Chem.* **1977**, *9*, 233–242.
- [24] a) H. Baumgarth, G. Meier, T. Braun, B. Braun-Cula, *Eur. J. Inorg. Chem.* **2016**, *2016*, 4565–4572; b) J. Fawcett, D. A. J. Harding, E. G. Hope, K. Singh, G. A. Solan, *Dalton Trans.* **2010**, *39*, 10781–10789; c) P. G. Watson, E. Lork, R. Mews, *J. Chem. Soc., Chem. Commun.* **1994**, 1069; d) E. A. V. Ebsworth, N. Robertson, L. J. Yellowlees, *J. Chem. Soc., Dalton Trans.* **1993**, 1031; e) S. A. Brewer, J. H. Holloway, E. G. Hope, P. G. Watson, *J. Chem. Soc., Chem. Commun.* **1992**, 1577–1578; f) A. J. Blake, R. W. Cockman, E. A. V. Ebsworth, J. H. Holloway, *J. Chem. Soc., Chem. Commun.* **1988**, 529.
- [25] E. A. V. Ebsworth, S. Moreton, *J. Organomet. Chem.* **1992**, *439*, 213–220.
- [26] A. Conkie, E. A. V. Ebsworth, R. A. Mayo, S. Moreton, *J. Chem. Soc., Dalton Trans.* **1992**, 2951.
- [27] a) T. Klapötke, *Polyhedron* **1989**, *8*, 311–315; b) M. F. A. Dove, J. C. P. Sanders, E. L. Jones, M. J. Parkin, *J. Chem. Soc., Chem. Commun.* **1984**, 1578–1581.
- [28] C. J. Bourgeois, S. A. Garratt, R. P. Hughes, R. B. Larichev, J. M. Smith, A. J. Ward, S. Willemsen, D. Zhang, A. G. DiPasquale, L. N. Zakharov, A. L. Rheingold, *Organometallics* **2006**, *25*, 3474–3480.
- [29] a) C. Röhr, S. Loss, *Z. Naturforsch. B* **1998**, *53*, 75–80; b) T. M. Klapötke, *J. Chem. Soc., Dalton Trans.* **1997**, 553–558; c) F. Jost, Y. Yacoby, D. Heitmann, S. Roth, *Phys. Rev. B* **1989**, *39*, 5444–5451; d) R. Mews, *Z. Naturforsch. B* **1973**, *28*, 99–100; e) L. C. Duncan, M. Kramar, *Inorg. Chem.* **1971**, *10*, 647–650.
- [30] a) M.-A. Guillevic, C. Rocaboy, A. M. Arif, I. T. Horváth, J. A. Gladysz, *Organometallics* **1998**, *17*, 707–717; b) G. Franciò, R. Scopelliti, C. G. Arena, G. Bruno, D. Drommi, F. Faraone, *Organometallics* **1998**, *17*, 338–347; c) R. D. Bedford, P. J. Dyson, A. F. Hill, A. G. Hulkes, C. J. Yates, *Organometallics* **1998**, *17*, 4117–4120; d) J. A. Cabeza, C. Landázuri, L. A. Oro, D. Belletti, A. Tiripicchio, M. T. Camellini, *J. Chem. Soc., Dalton Trans.* **1989**, 1093–1100; e) G. Yoneda, S.-M. Lin, L.-P. Wang, D. M. Blake, *J. Am. Chem. Soc.* **1981**, *103*, 5768–5771; f) E. A. V. Ebsworth, T. E. Fraser, *J. Chem. Soc., Dalton Trans.* **1979**, 1960–1964; g) C. Eaborn, N. Farrell, J. L. Murphy, A. Pidcock, *J. Chem. Soc., Dalton Trans.* **1976**, 58–67; h) M. A. Salomon, T. Braun, I. Krossing, *Dalton Trans.* **2008**, 5197–5206; i) P. Kläring, A.-K. Jungton, T. Braun, C. Müller, *Eur. J. Inorg. Chem.* **2012**, *2012*, 1430–1436; j) P. Kläring, S. Pahl, T. Braun, A. Penner, *Dalton Trans.* **2011**, *40*, 6785–6791.
- [31] a) O. Mahé, A. L'Heureux, M. Couturier, C. Bennett, S. Clayton, D. Tovell, F. Beaulieu, J.-F. Paquin, *J. Fluorine Chem.* **2013**, *153*, 57–60; b) A. Jesih, *J. Fluorine Chem.* **2000**, *103*, 25–29; c) L. N. Markovskii, V. E. Pashinnik, E. P. Saenko, *J. Org. Chem. USSR (Engl. Transl.)* **1977**, *13*, 1025–1026; *Zh. Organicheskoi. Khimii* **1977**, *13*, 1116–1117; d) L. N. Markovskii, L. S. Bobkova, V. E. Pashinnik, *J. Org. Chem. USSR (Engl. Transl.)* **1981**, *17*, 1699–1702; *Zh. Organicheskoi. Khimii* **1981**, *17*, 1903–1908.
- [32] G. M. Sheldrick, University of Göttingen, Germany, **1996**.
- [33] a) G. M. Sheldrick, *Acta Crystallogr., Sect. A* **2015**, *71*, 3–8; b) G. M. Sheldrick, *Acta Crystallogr., Sect. C* **2015**, *71*, 3–8.

Received: January 23, 2020

Published Online: May 19, 2020